

Accelerating Cosmologies and a Phase Transition in M-Theory

Mattias N.R. Wohlfarth*

*Department of Applied Mathematics and Theoretical Physics,
Centre for Mathematical Sciences, University of Cambridge,
Wilberforce Road, Cambridge CB3 0WA, U.K.*

(Dated: 1st February 2008)

Abstract

M-theory compactifies on a seven-dimensional time-dependent hyperbolic or flat space to a four-dimensional FLRW cosmology undergoing a period of accelerated expansion in Einstein conformal frame. The strong energy condition is violated by the scalar fields produced in the compactification, as is necessary to evade the no-go theorem for time-independent compactifications. The four-form field strength of eleven-dimensional supergravity smoothly switches on during the period of accelerated expansion in hyperbolic compactifications, whereas in flat compactifications, the three-form potential smoothly changes its sign. For small acceleration times, this behaviour is like a phase transition of the three-form potential, during which the cosmological scale factor approximately doubles.

PACS numbers: 11.25.Yb, 98.80.Jk, 04.50.+h

At present, the study of accelerating cosmologies deserves particular interest for several reasons. Recent astronomical measurements on certain types of supernovae, as reported, e.g., in [1], seem to indicate that our universe is not only expanding, which has been known for a long time, but is undergoing an accelerated expansion. Moreover, the hypothesis of another, inflationary, epoch of accelerated expansion in the distant past is supported by measurements on the cosmic microwave background, see, e.g., [2]. Phenomenologically motivated cosmological models which exhibit accelerated expansion for at least a certain period of time are easily constructed. The task, however, of deriving such a model from a compactification of a higher-dimensional theory such as superstring or M-theory, which are believed to incorporate the standard model consistently, is a more difficult one. This is due to the existence of a no-go theorem for compactifications on *time-independent* internal spaces [3, 4]. This theorem states that if the strong energy condition is satisfied for a higher-dimensional theory, which it is for ten or eleven-dimensional string or M-theory, then it is also valid for the compactified theory. But then the time-time component of the four-dimensional Ricci tensor is subject to the condition $R_{00} \geq 0$, and, since R_{00} is related to the acceleration of the scale factor S in an FLRW universe by $R_{00} = -3\ddot{S}/S$, it follows that acceleration is forbidden.

A recent letter demonstrated the possibility of evading this theorem [5]; a solution of the vacuum Einstein equations was presented for which spacetime is compactified on a compact hyperbolic manifold of *time-varying* volume to a four-dimensional homogeneous and isotropic FLRW universe with a phase of accelerated expansion. This work is easily extended to the M-theory case where, in the low energy limit, the equations of motion follow from the bosonic part of the action of eleven-dimensional supergravity [6],

$$S = \int_{11} \sqrt{-g} \left(R - \frac{1}{2 \cdot 4!} F_4^2 \right), \quad (1)$$

involving the curvature scalar R and the four-form field strength $F_4 = dC_3$. In principle, this action should additionally contain the Chern-Simons term $C_3 \wedge F_4 \wedge F_4$. This term, however, becomes irrelevant below when F_4 is realized as a volume form. The fact that F_4 can be non-zero in toroidal or spherical compactifications has been established before [7, 8]. One new point here is to look at the remaining possibility and to investigate the effects of a non-zero F_4 in a hyperbolic compactification.

Consider the following eleven-metric, parameterized by functions of time $K(t)$, $L(t)$ and

$S(t)$,

$$ds_{11}^2 = \left(\frac{K(t)}{L(t)} \right)^{-7/6} ds_E^2 + \left(\frac{K(t)}{L(t)} \right)^{1/3} ds_7^2, \quad (2)$$

where

$$ds_E^2 = -S(t)^6 dt^2 + S(t)^2 d\mathbf{x} \cdot d\mathbf{x} \quad (3)$$

is a four-dimensional FLRW metric in Einstein conformal frame. The internal space with the metric

$$ds_7^2 = g_{7mn} dy^m dy^n \quad (4)$$

is taken to be a compact Einstein manifold with negative or zero curvature, such that its Ricci tensor satisfies

$$R_{7mn} = -6\kappa^2 g_{7mn} \quad \text{for } \kappa \geq 0. \quad (5)$$

The eleven-metric (2), together with the four-form

$$F_4 = bL(t)^2 vol_{1,3} \quad (6)$$

where $vol_{1,3} = dt \wedge dx_1 \wedge dx_2 \wedge dx_3$, solves the Einstein equations and the equation of motion of F_4 derived from the action (1) if

$$S(t) = K(t)^{7/12} L(t)^{-1/4} \quad (7)$$

and

$$K(t) = \begin{cases} \frac{\sqrt{21}\gamma}{14\kappa \sinh \frac{3}{7}\sqrt{21}\gamma|t|} & \text{for } \kappa > 0 \\ e^{\frac{3}{7}\sqrt{21}\gamma t} & \text{for } \kappa = 0 \end{cases}, \quad (8a)$$

$$L(t) = \frac{3\gamma}{b \cosh 3\gamma t}. \quad (8b)$$

Note that not all constants of integration are displayed: different time shifts may be chosen in $K(t)$ and $L(t)$. Solutions that are of similar type have been considered before, however, without appreciating their possible accelerating behaviour, see e.g. [9]. Moreover, the specific solution presented above has a close relation to S -branes, see [10, 11] (and references therein). It describes, in fact, an electrically charged $SM2$ -brane, with a hyperbolic or flat transverse space, whose three spatial worldvolume directions correspond to the spatial sections of the four-dimensional FLRW cosmology considered above. It is clear from $K(t)$ in (8) that the $\kappa > 0$ solution with hyperbolic internal space does not simply reduce to the




Figure 1: Plot of a typical scale factor $S(\eta)$ and its acceleration $d^2S/d\eta^2$ (multiplied by S^5).

one for $\kappa = 0$ where the internal space is flat. The same happens for the field strength: the solution for $L(t)$ with non-zero field strength parameter $b \neq 0$ does not simply reduce to the one for $b = 0$, given in [5], where instead $L(t) = e^{3\gamma t}$ (with γ set to $\gamma = 1$).

The standard time coordinate η in Einstein frame is defined by $d\eta = S(t)^3 dt$, see (3). Expansion of the Einstein frame scale factor corresponds to $dS/d\eta > 0$ and acceleration to $d^2S/d\eta^2 > 0$. Hence, as in the zero field strength case [5], it is easily shown that the above solutions admit a period of accelerated expansion for negative t in a certain interval

$$\frac{t_1}{\gamma} < t < \frac{t_2}{\gamma} \quad (9)$$

(which maps into an interval $\eta_1 < \eta < \eta_2$ of positive η), see figure 1. Such accelerated expansion, however, can only be possible if the strong energy condition is violated by the scalar fields arising in the compactification process. These are the scale e^M of the internal space, which is determined by

$$M(t) = \frac{1}{6} \ln \frac{K(t)}{L(t)}, \quad (10)$$

and a term coming from F_4 . Reducing the eleven-dimensional action (1) by substituting the metric ansatz (2), one finds a four-dimensional scalar Lagrangian

$$\mathcal{L} = 7\nabla_E^2 M - \frac{63}{2}(\nabla_E M)^2 - 42\kappa^2 e^{-9M} - \frac{1}{2 \cdot 4!} e^{21M} F_4^2. \quad (11)$$

Note that this expression is given in Einstein frame and is only reached by a suitable conformal transformation of the four-dimensional part of the original eleven-metric. It is necessary to go to the Einstein frame in order to avoid interpretational ambiguities: in particular, this

frame is characterized by a time-independent Newton constant. From the energy-momentum tensor $T_{\mu\nu}$ of $\mathcal{L}$ one finds

$$\begin{aligned} X &\equiv T_{\eta\eta} - \frac{1}{2}Tg_{E\eta\eta} \\ &= 63M'^2 - 42\kappa^2 e^{-9M} - \frac{1}{2}b^2 e^{-21M} \end{aligned} \quad (12)$$

where $M' = dM/d\eta$. This quantity is required to be positive by the strong energy condition, and indeed,

$$X = -6S^{-1} \frac{d^2 S}{d\eta^2}, \quad (13)$$

such that the strong energy condition is violated if, and only if, the acceleration of the scale factor S is positive. Note, from the second term in X , that a hyperbolic internal space, where $\kappa > 0$, gives rise to a negative contribution. For a flat internal space, this term vanishes and the strong energy condition can be violated only by the third term which is due to the four-form F_4 . The latter contributes negatively because of a more general observation [3]: p -forms in n dimensions may violate the strong energy condition for $p \geq n - 1$. Such an effect, that the four-form of eleven-dimensional supergravity gives rise to a negative cosmological constant upon reduction to four dimensions, has also been noted in [7].

The phase of acceleration has several interesting properties. In the following they are discussed for a hyperbolic internal space. For a flat internal space one proceeds similarly, and a summary of the results for this case is given below. Now let $\kappa > 0$. Accelerated expansion starts at the time t_1/γ and ends at t_2/γ where one finds $t_1 \cong -0.78$ and $t_2 \cong -0.15$ as the two negative solutions of the equation

$$\begin{aligned} &-(5 - \sqrt{21}) \cosh 6(1 + \frac{1}{7}\sqrt{21})t + 8 \cosh \frac{6}{7}\sqrt{21}t \\ &-(5 + \sqrt{21}) \cosh 6(1 - \frac{1}{7}\sqrt{21})t + 8 \cosh 6t - 10 = 0 \end{aligned} \quad (14)$$

which follows from $d^2 S/d\eta^2 = 0$. In terms of the physical time variable η , the accelerating phase is bounded by

$$\eta_i = A(t_i) b^{3/4} \kappa^{-7/4} \quad \text{for } i = 1, 2 \quad (15)$$

where the function $A(t)$ is given by

$$A(t) = \left(\frac{3}{2^{14} 7^7} \right)^{1/8} \int_{-\infty}^t ds \frac{\cosh^{3/4} 3s}{\sinh^{7/4} \frac{3}{7}\sqrt{21}|s|}. \quad (16)$$




Figure 2: *The scale of the internal space given by $M(\eta)$.*

The difference $\eta_{\text{acc}} = \eta_2 - \eta_1$ gives the total acceleration time

$$\eta_{\text{acc}} = [A(t_2) - A(t_1)]b^{3/4}\kappa^{-7/4}, \quad (17)$$

and the expansion factor during the period of accelerated expansion is obtained to be

$$a = \frac{S(\eta_2)}{S(\eta_1)} \cong 2.155. \quad (18)$$

Note that the factor a is constant and independent of the parameters of the solution (8). This implies that there can be no hope of using this solution in any inflationary scenario, where the required expansion to solve cosmological issues, such as the horizon or the flatness problem, is of the order of magnitude $\mathcal{O}(e^{60})$. The total acceleration time η_{acc} , on the other hand, depends on the field strength parameter b as well as on the curvature κ of the internal space. It tends to zero for $b \rightarrow 0$ or $\kappa \rightarrow \infty$ in which limit the scale factor doubles instantaneously. The scale of the internal space is determined by the behaviour of M in (10). One finds that $M \rightarrow \infty$ as $\eta \rightarrow 0$ and also as $\eta \rightarrow \infty$. In the accelerating phase, M assumes a minimum value, as is shown in figure 2. Looking at the solution in the whole range of η shows that there is no stabilisation of the internal space scale, which seems to be a more general problem in hyperbolic compactifications [12]. The fact, however, that the scale has a minimum, which is reached very fast, means that there is a natural ‘compactification’ of the internal space until the end of the accelerating phase. The following decompactification is a very slow process.




Figure 3: *The behaviour of the four-form F_4 as a function of η (for a hyperbolic internal space $\kappa > 0$).*

The four-form field F_4 smoothly switches on in the phase of accelerated expansion. For small times $\eta \rightarrow 0$, it behaves like

$$F_4 \stackrel{\eta \rightarrow 0}{\sim} b^{-(1+2\epsilon)} \gamma^2 \kappa^{14\epsilon/3} \eta^{8\epsilon/3} \text{vol}_{1,3} \quad (19)$$

where $\epsilon = (\sqrt{21}/3 - 1)^{-1}$, whereas in the limit $\eta \rightarrow \infty$, the field F_4 tends to a constant value $F_4 = F_\infty \text{vol}_{1,3}$ with

$$F_\infty = 9\gamma^2/b. \quad (20)$$

More precisely,

$$F_4 \stackrel{\eta \rightarrow \infty}{\sim} \left(F_\infty - 9\gamma^2 b (6\kappa)^{-14/3} \left[\frac{3}{4} (\eta - \eta_0) \right]^{-8/3} \right) \text{vol}_{1,3}, \quad (21)$$

see also figure 3. As $b \rightarrow 0$, the four-form value at infinity F_∞ may be kept constant by choosing an appropriate value for γ . But this limit coincides with the one of zero total acceleration time. For small b , therefore, the three-form potential C_3 of eleven-dimensional supergravity is behaving as in a first order phase transition, characterized by the fact that its derivative $F_4 = dC_3$ becomes discontinuous. The four-form field F_4 is almost instantaneously switched on to its $\eta \rightarrow \infty$ value $F_\infty \text{vol}_{1,3}$ while the scale factor $S(\eta)$ of the FLRW universe jumps by the factor $a \cong 2.155$.

Many of the results for a flat internal space, where $\kappa = 0$, are similar to those for the hyperbolic case $\kappa > 0$, some are very different. The appropriate expressions (15, 17) for




Figure 4: *The behaviour of the four-form F_4 as a function of η (for a flat internal space $\kappa = 0$).*

the times which characterize the period of accelerated expansion are essentially obtained by replacing κ by γ (the integral in (16) differs, of course). The expansion factor of the FLRW cosmology in this period becomes $a \cong 1.355$. For the internal scale one again finds an evolution as shown in fig. 2. The interesting changes appear in the behaviour of F_4 . While (19) stays the same up to the replacement $\kappa \mapsto \gamma$, one finds that F_4 tends to zero also for $\eta \rightarrow \infty$. For some finite $\eta_{\max}$, the four-form assumes a maximum value which equals the limiting value of the $\kappa > 0$ case, $F_{\max} = F_{\infty}$. For a plot of F_4 see figure 4. These differences between $\kappa > 0$ and $\kappa = 0$ are best understood from the three-form potential

$$C_3 = \frac{3\gamma}{b} \tanh 3\gamma t \, dx_1 \wedge dx_2 \wedge dx_3. \quad (22)$$

In the hyperbolic case $\kappa > 0$, a negative C_3 at $\eta = 0$ ($t = 0$) is switched off as $\eta \rightarrow \infty$ ($t \rightarrow 0$). The value F_{∞} of F_4 is the slope at $t \rightarrow 0$. The difference for $\kappa = 0$ is that, here, $\eta \rightarrow \infty$ corresponds to $t \rightarrow \infty$ because $S(t)$ is non-singular. Thus, a negative C_3 at $\eta = 0$ ($t = 0$) makes a transition to a positive C_3 of the same absolute value as $\eta \rightarrow \infty$ ($t \rightarrow \infty$). This transition takes place around $\eta_{\max}$. It has the slope $F_{\max}$ and can be made discontinuous in the limit $b \rightarrow 0$ while keeping the ratio γ/b constant.

The alignment of the four-form along the directions (0 1 2 3) of the final cosmology is not the only possibility consistent with the rotational symmetry of FLRW spacetimes. Rather than choosing F_4 as above, one might also consider aligning it along (0 | 4 5 6), (1 2 3 | 4) or along directions (4 5 6 7) completely in the internal space. Another interesting question is whether there are solutions where the Chern-Simons term of the eleven-dimensional su-

pergravity action becomes important, or whether it is possible to find solutions where the internal space is a product of several manifolds. With such extensions, one might hope to be able to control the properties of the phase of accelerated expansion or even to combine several of these phases into a single solution. This then might provide a key to derive more realistic cosmologies, including inflation, from M-theory. After completion of this work, several papers have appeared which explore further aspects of accelerating cosmologies from string and M-theory compactifications [13, 14, 15].

Acknowledgments. MNRW thanks Paul K. Townsend for valuable discussions. He gratefully acknowledges financial support from the Gates Cambridge Trust.

* Electronic address: M.N.R.Wohlfarth@damtp.cam.ac.uk

- [1] A.G. Riess *et al.*, *The farthest known supernova: support for an accelerating universe and a glimpse of the epoch of deceleration*, *Astrophys. J.* **560**, 49 (2001).
- [2] C.L. Bennett *et al.*, *First year Wilkinson Microwave Anisotropy Probe (WMAP) observations: preliminary maps and basic results*, [[astro-ph/0302207](#)].
- [3] G.W. Gibbons, *Aspects of supergravity theories in Supersymmetry, supergravity and related topics*, edited by F. del Aguila, J.A. de Azcárraga and L. Ibáñez, 123 (World Scientific, Singapore, 1985).
- [4] J. Maldacena and C. Nuñez, *Supergravity description of field theories on curved manifolds, and a no go theorem*, *Int. J. Mod. Phys.* **A16**, 822 (2001).
- [5] P.K. Townsend and M.N.R. Wohlfarth, *Accelerating cosmologies from compactification*, submitted to *Phys. Rev. Lett.*, [[hep-th/0303097](#)].
- [6] E. Cremmer, B. Julia and J. Scherk, *Supergravity theory in eleven dimensions*, *Phys. Lett.* **B76**, 409 (1978).
- [7] A. Aurilia, H. Nicolai and P.K. Townsend, *Hidden constants: the θ parameter of QCD and the cosmological constant of $N = 8$ supergravity*, *Nucl. Phys.* **B176**, 509 (1980).
- [8] P.G.O. Freund and M.A. Rubin, *Dynamics of dimensional reduction*, *Phys. Lett.* **B97**, 233 (1980).
- [9] J. Demaret, J.-L. Hanquin, M. Henneaux and P. Spindel, *Cosmological models in eleven-dimensional supergravity*, *Nucl. Phys.* **B252**, 538 (1985).

- [10] N. Ohta, *Intersection rules for S-branes*, Phys. Lett. **B558**, 213 (2003); *Accelerating cosmologies from S-branes*, [[hep-th/0303238](#)].
- [11] S. Roy, *Accelerating cosmologies from M/string theory compactifications*, [[hep-th/0304084](#)].
- [12] S. Nasri, P.J. Silva, G.D. Starkman, and M. Trodden, *Radion stabilization in compact hyperbolic extra dimensions*, Phys. Rev. **D66**, 045029 (2002).
- [13] R. Emparan and J. Garriga, *A note on accelerating cosmologies from compactifications and S-branes*, [[hep-th/0304124](#)].
- [14] N. Ohta, *A study of accelerating cosmologies from superstring/M theories*, [[hep-th/0304172](#)].
- [15] C.-M. Chen, P.-M. Ho, I.P. Neupane and J.E. Wang, *A note on acceleration from product space compactification*, [[hep-th/0304177](#)].